

Newly Report of the First Zoea of *Zehntneriana amakusae* (Brachyura: Pilumnidae: Rhizopinae) Hatched in Laboratory Material from Korea

Seok Hyun Lee*

Department of Companion Animals, Silla University, Busan 46958, Korea

ABSTRACT

Ovigerous crab of *Zehntneriana amakusae* of Rhizopinae in Pilumnidae was collected from Munseom Islet, Seogwipo, Jeju Is. and the first zoeas were obtained from hatched materials in the laboratory. The morphology of the first zoea of *Z. amakusae* is reported for the first time in the world with a color digital photo. The first zoea of *Z. amakusae* has common zoeal characteristics in the spines of carapace and the saturations of maxillule, maxilla, and the first maxilliped of Korean Pilumnids. However, the first zoea of *Z. amakusae* is different from other species of Korean Pilumnids in that it has 1, 1, 5 setae on endopod of the second maxilliped and two lateral spines and one dorsal spine on fork of telson.

Keywords: *Zehntneriana amakusae*, first zoea, Rhizopinae, Pilumnidae, Korea

INTRODUCTION

The life cycle of brachyuran crabs generally consists of naupliar (passed in egg), zoeal (planktonic life, consisting of at least two stages), megalopal (introducing benthic life, one or two stages), and adult (within benthic life) stages (Hong, 2021). To date, most studies of the classification and phylogeny systematics of decapods have been relied on adult morphologies, however, it is difficult to study using the adult morphology with many variations because they are adapted to various habitats (Souza et al., 2013). Most larvae inhabit a planktonic environment, exhibiting fewer morphological variations than adult and aligning more closely with molecular data (Rice, 1980; Hultgren et al., 2009). Therefore, extensive efforts are still required to describe brachyuran larval morphologies.

Currently, 15 species of three subfamilies of the family Pilumnidae have been reported from Korean waters (Lee et al., 2021, 2023). Among them, nine species of zoeal morphologies are known in the Korea (Table 1). The genus *Zehntneriana* Ng and Takeda, 2010 of the subfamily Rhizopinae Stimpson, 1858 includes five species [*Zehntneriana amakusae* (Takeda and Miyake, 1969); *Z. miyakei* (Takeda, 1972); *Z. novaeinsulicola* (Takeda and Kurata, 1977); *Z. serrata* Ng and Lin, 2015; *Z. tadafumii* Lee, Kim and Ng, 2015; *Z. villosa*

(Zehntner, 1894)] on record in the world (WoRMS, 2024). Among them, larvae have not been reported yet. During the study about decapod larvae collected an ovigerous crab, and identified as *Z. amakusae* by having covered dense short setae on frontal, supraorbital, and anterolateral regions of transversely ovoid carapace, and divided by three crest-shaped teeth on anterolateral margin. The first zoeas of *Z. amakusae* were obtained from the laboratory hatching materials for the first time in the world.

The purposes of the present study are to illustrate and describe the morphology of the first zoea of *Z. amakusae* for the first time in the world, providing a color digital photo, and comparing its morphological characteristics with those of other the first zoeas of the family Pilumnidae.

MATERIALS AND METHODS

An ovigerous crab of *Zehntneriana amakusae* was collected by SCUBA diving at a depth of approximately 20 m in Munseom Islet, Seogwipo, Jeju Island, Republic of Korea on 13 Aug 2010 and its zoea hatched in the laboratory on 22 Aug 2010. The first zoeas were first fixed in 30% ethanol and then preserved in 94% ethanol for examination. A color digital photo

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

***To whom correspondence should be addressed**
Tel: 82-51-999-6931, Fax: 82-51-999-5471
E-mail: sky3284@silla.ac.kr

Table 1. List of species of the family Pilumnidae from Korean waters and its known larval stages

Family	Subfamily	Genus	Species	Larval stages	References	
Pilumnidae	Eumedoninae	<i>Echinoecus</i>	<i>nipponicus</i>	Z1	Lee (2020)	
		Pilumninae	<i>Actumnus</i>	<i>elegans</i> <i>marissinicus</i>		
	<i>Benthopanope</i>		<i>indica</i>	Z1-4, M	Ko (1995)	
	<i>Harrovia</i>		<i>elegans</i> <i>japonica</i>	Z1-4, M	Lee and Ko (2009)	
	<i>Heteropilumnus</i>		<i>ciliatus</i>	Z1-3	Ko and Yang (2003)	
	<i>Neoactumnus</i>		<i>convexus</i>			
	<i>Pilumnopeus</i>		<i>granulatus</i> <i>makianus</i>	Z1-4, M Z1-4	Ko (1997) Lee (1993)	
	Rhizopinae		<i>Arges</i> <i>Zehntneriana</i>	<i>longicornis</i>	Z1	Clark and Paula (2003)
				<i>minutus</i>	Z1-4, M	Ko (1994a, 1997)
				<i>trispinosus</i>	Z1-4, M	Ko (1994b)
				<i>parallelus</i>		
				<i>amakusae</i>	Z1	Present study

Z, zoeal stage; M, megalopal stage.



Fig. 1. Color digital photos of adult and the first zoeas of *Zehntneriana amakusae*. A, Adult of *Z. amakusae*; B, First zoeas of *Z. amakusae*.

of living zoeas for observation of chromatophore patterns was taken with Leica EZ40 microscope (Wetzlar, Germany) and subsequently processed in Photoshop (Photoshop 2021, Adobe). The dissection and appendage examined of zoeal specimens and drawings of the morphology of zoea followed the described by Lee (2020).

The sequence of zoeal description follows that of Clark and Cuesta (2015). The observation of chromatophore patterns was determined through a color digital photo of living zoeas. The measurements of zoeas were made using a micrometer. The carapace length and rostral and dorsal carapace spines length were abbreviated as CL and RDL, and their meanings are described in Lee (2020). The examined zoeal specimens and

spent female crab have been deposited at the Silla University, Korea.

RESULTS

Order Decapoda Latreille, 1802
Family Pilumnidae Samouelle, 1819
Subfamily Rhizopinae Stimpson, 1858
Genus *Zehntneriana* Ng and Takeda, 2010

Zehntneriana amakusae (Takeda & Miyake, 1969)

Description of Zoea I (Figs. 1B, 2). Size: CL 0.51 ± 0.02




Fig. 2. *Zehntneriana amakusae*, first zoea. A, Lateral view; A', Lateral expansion of carapace; B, Antennule; C, Antenna; D, Mandibles; E, Maxillule; F, Maxilla; G, First maxilliped; H, Second maxilliped; I, Dorsal view of pleon and telson; I', Fork of telson. Scale bars: A, I=0.2 mm, A'-H, I'=0.1 mm.

Table 2. Comparison of the characteristics of the first zoeae of family Pilumnidae in Korean waters

	Pilumnidae									
	Eumedininae			Pilumninae				Rhizopinae		
Subfamily	<i>Echinoecus nipponicus</i>			<i>Pilumnopus granulata</i>				<i>Zehntneriana amakusae</i>		
Species	<i>Harrovia japonica</i>	<i>Benthopanope indica</i>	<i>Heteropilumnus ciliatis</i>	<i>Pilumnopus granulata</i>	<i>P. makiana</i>	<i>Pilumnus longicornis</i>	<i>P. minutus</i>	<i>P. trispinosus</i>		
Reference	Lee and Ko (2009)	Ko (1995)	Ko and Yang (2003)	Ko (1997)	Lee (1993)	Clark and Paula (2003)	Ko (1994a)	Ko (1994b)	Present study	
Carapace										
Dorsal, rostral, lateral spines	Long, short, short	Long, short, absent	Long, long, short	Long, vestigial, short	Long, long, short	Long, long, short	Long, short, short	Long, short, absent	Long, short, short	
Antennule	3 aesthetascs, 2 setae	3 aesthetascs, 2 setae	4 aesthetascs, 2 setae	3 aesthetascs, 3 setae	4 aesthetascs, 1 seta	4 aesthetascs, 2 setae	3 aesthetascs, 3 setae	3 aesthetascs, 3 setae	3 aesthetascs, 2 setae	
Antenna										
Protopod	Spinulate	Spinulate	Spinulate	Spinulate	Spinulate	Spinulate	Spinulate	Spinulate	Spinulate	
Exopod	Shorter than protopod, spinulate	Longer than protopod, spinulate	Longer than protopod, spinulate	As same as protopod, spinulate	Longer than protopod, spinulate	As same as protopod, spinulate	Shorter than protopod, spinulate	Longer than protopod, spinulate	Longer than protopod, spinulate	
Medial setae	2 unequal	2 unequal	2 unequal	2 unequal	2 unequal	2 unequal	2 unequal	2 unequal	2 unequal	
Maxillule										
Coxal endite	7 setae	7 setae	7 setae	7 setae	5 setae	7 setae	7 setae	7 setae	7 setae	
Basal endite	5 setae	5 setae	5 setae	5 setae	5 setae	5 setae	5 setae	5 setae	5 setae	
Endopod	1, 2+4 setae	1, 2+4 setae	1, 2+4 setae	1, 2+4 setae	1, 2+4 setae	1, 2+4 setae	1, 2+4 setae	1, 2+4 setae	1, 2+4 setae	
Maxilla										
Coxal endite	6+4 setae	6+4 setae	6+4 setae	6+4 setae	6+3 setae	6+4 setae	6+4 setae	6+4 setae	6+4 setae	
Basal endite	5+4 setae	5+4 setae	5+4 setae	5+4 setae	5+3 setae	5+4 setae	5+4 setae	4+5 setae	4+5 setae	
Endopod	3+5 setae	3+5 setae	3+5 setae	3+5 setae	3+5 setae	3+5 setae	3+5 setae	3+5 setae	3+5 setae	
1st maxilliped										
Basis	2, 2, 3, 3 setae	2, 2, 3, 3 setae	2, 2, 3, 3 setae	2, 2, 3, 3 setae	2, 2, 3, 3 setae	2, 2, 3, 3 setae	2, 2, 3, 3 setae	2, 2, 3, 3 setae	2, 2, 3, 3 setae	
Endopod	3, 2, 1, 2, 5 setae	3, 2, 1, 2, 5 setae	3, 2, 1, 2, 5 setae	3, 2, 1, 2, 5 setae	3, 2, 1, 2, 5 setae	3, 2, 1, 2, 5 setae	3, 2, 1, 2, 5 setae	3, 2, 1, 2, 5 setae	3, 2, 1, 2, 5 setae	
2nd maxilliped										
Basis	1, 1, 1, 1 setae	1, 1, 1, 1 setae	1, 1, 1, 1 setae	1, 1, 1, 1 setae	1, 1, 1, 1 setae	1, 1, 1, 1 setae	1, 1, 1, 1 setae	1, 1, 1, 1 setae	1, 1, 1, 1 setae	
Endopod	1, 1, 6 setae	1, 1, 6 setae	1, 1, 6 setae	1, 1, 6 setae	1, 1, 6 setae	1, 1, 6 setae	1, 1, 6 setae	1, 1, 6 setae	1, 1, 6 setae	
Pleon										
Lateral processes	Pleomeres 2, 3	Pleomere 2	Pleomeres 2, 3	Pleomeres 2, 3	Pleomeres 2, 3	Pleomeres 2, 3	Pleomeres 2-5	Pleomeres 2-5	Pleomeres 2, 3	
Telson										
Fork	2 lateral spines, 1 dorsomedial spine	1 lateral spine, 1 lateral seta, 1 dorsomedial spine	1 lateral spine, 1 lateral seta, 1 dorsomedial spine	1 lateral spine, 1 lateral seta, 1 dorsomedial spine	1 lateral spine, 1 lateral seta, 1 dorsomedial spine	2 lateral spines, 1 dorsomedial spine	1 lateral spine, 1 lateral seta, 1 dorsomedial spine	1 lateral spine, 1 lateral seta, 1 dorsomedial spine	1 lateral spine, 1 lateral seta, 1 dorsomedial spine	
Spinulate	Spinulate	Spinulate	Spinulate	Spinulate	Spinulate	Spinulate	Spinulate	Spinulate	Spinulate	

mm; RDL 1.03 ± 0.02 mm.

Chromatophores (Fig. 1B): Yellowish brown chromatophores which occurring behind eyes, on basis of dorsal spine posteriorly, on basal of basis of each maxillipeds, on abdominal somites 1, 2 ventrally and margins of abdominal somites 4, 5 posteriorly.

Carapace (Figs. 1B, 2A, A'): dorsal spine curved distally, approximately equal CL, with several minute tubercles; rostral spine shorter than antennal protopod length, without digital spinulation; lateral spines short, curved; anterodorsal spine absent; 1 pair of posterodorsal setae present; ventral margin with spinules.

Cephalon. Eyes (Figs. 1B, 2A): sessile.

Antennule (Fig. 2B): primary flagellum unsegmented with 3 (2 broad, 1 slender) aesthetascs, of unequal length plus 2 terminal setae of unequal length; accessory flagellum absent.

Antenna (Fig. 2C): uniramous; protopod distal medially bilaterally spinulate, longer than rostral spine; endopod bud absent; exopod distal medially bilaterally spinulate with 2 medial setae of unequal length.

Mandibles (Fig. 2D): asymmetrical; right molar with 4 teeth, left molar with 1 tooth, confluent with incisor process; palp absent.

Maxillule (Fig. 2E): uniramous; epipod seta absent; coxal endite with 7 setae; basal endite with 5 terminal setal processes + 2 small setal buds; endopod comprising 2 articles, proximal article with 1 seta, distal article with 6 (2 subterminal + 4 terminal) setae; exopod seta absent.

Maxilla (Fig. 2F): Coxal endite bilobed, with 6 + 4 setae; basal endite bilobed, with 5 + 4 setae; endopod bilobed, with 3 + 2 + 3 setae; exopod (scaphognathite) margin with 4 plumose setae, 1 distal process.

Pereion. First maxilliped (Fig. 2G): biramous; coxa without setae; basis with 10 (2 + 2 + 3 + 3) setae; endopod comprising 5 articles with 3, 2, 1, 2, 5 (1 subterminal + 4 terminal) setae respectively; exopod comprising 2 articles, distal article with 4 long terminal plumose natatory setae.

Second maxilliped (Fig. 2H): biramous; coxa without setae; basis with 4 (1 + 1 + 1 + 1) setae; endopod comprising 3 articles, with 1, 1, 5 (2 subterminal + 3 terminal) setae; exopod comprising 2 articles, distal article with 4 long terminal plumose natatory setae.

Third maxilliped: absent.

Pereiopods: absent.

Pleon (Fig. 2I): 5 pleomeres; pleomere 2 with pair of lateral processes directed laterally; pleomere 3 with pair of lateral processes directed posteriorly; pleomeres 2–5 each with 1 pair of posterodorsal setae and spinulated dorsal posterior margin; pleopod bud absent.

Telson (Fig. 2I'): each fork long, spinulated, margin with 1 long lateral spine, 1 small lateral spine, 1 stout dorsomedial

spine present; posterior margin with 3 pairs of stout spinulate setae.

DISCUSSION

The first zoea of *Zehntneriana amakusae* is illustrated and described for the first time in the world with a color digital photo. The first zoea of *Z. amakusae* has yellowish brown chromatophores located behind the eyes, on the basis of dorsal spine posteriorly, on the basal of basis of each maxillipeds, on pleomeres 1, 2 ventrally and margins of pleomeres 4, 5 posteriorly, dorsal spine subequal carapace length, rostral, and lateral spines shorter than carapace length, three aesthetascs and two setae on the antennule, two medial setae on the antennal exopod, lateral processes on the pleomeres 2, 3, and two lateral spines and one dorsomedial spine on the telson.

Including present study, the first zoeal stages of 10 species of the family Pilumnidae have been reported in Korean waters (Tables 1, 2). Table 2 summarizes the characteristics of the first zoeal stage of the family Pilumnidae known in Korean waters. The first zoeas of Korean Pilumnids identically by having exopod of antenna with two unequal sized medial spine or seta, endopods of maxillule and maxilla with 1, 2 + 4 and 3 + 5 setae, respectively, and spinulated fork of telson with 2 lateral spines or setae and a dorsomedial spine as suggested by Ko and Lee (2012) (Table 2).

The zoeal characteristics of the first zoea of *Z. amakusae* are almost identical to other species belonging to the family Pilumnidae. However, it has slightly differed from the others by having an endopod of the second maxilliped with 1, 1, 5 setae and the fork of telson with two lateral spines and one dorsomedial spine (Table 2).

ORCID

Seok-Hyun Lee: <https://orcid.org/0000-0003-4372-4909>

CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

ACKNOWLEDGMENTS

This work was supported by the management of Marine Fishery Bio-resources Center (2024) funded by the National Marine Biodiversity Institute of Korea (MABIK).

REFERENCES

- Clark PF, Cuesta JA, 2015. Chapter 71-17. Larval systematics of Brachyura. In: Decapoda: Brachyura, treatise on zoology - anatomy, taxonomy, biology. The Crustacea, Vol. 9. Part C (2 volumes). Complementary to the volumes translated from the French of the *Traité de Zoologie* (Eds., Castro P, Davie PJF, Guinot D, Schram FR and von Vaupel Klein JC). Brill, Leiden and Boston, pp. 981-1048. https://doi.org/10.1163/9789004190832_020
- Clark PF, Paula J, 2003. Descriptions of ten xanthoidean (Crustacea: Decapoda: Brachyura) first stage zoeas from Inhaca Island, Mozambique. *The Raffles Bulletin of Zoology*, 51:323-378. <https://doi.org/10.5281/zenodo.13229521>
- Hong SY, 2021. Diversity of Crustacean Larvae. *Journal of National Academy of Science, Republic of Korea: Natural Science Series*, 60:279-316.
- Hultgren KM, Guerao G, Marques FPL, Palero FP, 2009. Assessing the contribution of molecular and larval morphological characters in a combined phylogenetic analysis of the superfamily Majoidea. In: *Decapod crustacean phylogenetics, crustacean issues 18* (Eds., Martin JW, Crandall KA, Felder DL). CRC Press, Boca Raton, FL, pp. 437-455.
- Ko HS, 1994a. The zoeal stages of *Pilumnus minutus* De Haan, 1835 (Decapoda: Brachyura: Pilumnidae) in the laboratory. *Korean Journal of Systematic Zoology*, 10:145-155.
- Ko HS, 1994b. Larval development of *Parapilumnus trispinosus* Sakai, 1965 (Crustacea, Brachyura, Xanthidae) reared in the laboratory. *Korean Journal of Zoology*, 37:331-342.
- Ko HS, 1995. Larval development of *Benthopanope indica* (De Man, 1887) (Decapoda: Brachyura: Pilumnidae) in the laboratory. *Journal of Crustacean Biology*, 15:280-290. <https://doi.org/10.2307/1548955>
- Ko HS, 1997. Larval development of *Pilumnopeus granulata* Balss, 1933 and *Pilumnus minutus* De Haan, 1835 (Crustacea: Brachyura: Pilumnidae), with a key to the known pilumnid zoeae. *Korean Journal of Biological Sciences*, 1:31-42. <https://doi.org/10.1080/12265071.1997.9647346>
- Ko HS, Lee SH, 2012. Invertebrate fauna of Korea. Vol. 21, No. 22. Crabs and zoeas II. Arthropoda: Malacostraca: Decapoda: Brachyura: Eriphioidea, Pilumnoidea, Xanthoidea. *National Institute of Biological Resources, Incheon*, pp. 1-93.
- Ko HS, Yang HJ, 2003. Zoeal development of *Heteropilumnus ciliatus* (Decapoda: Pilumnidae), with a key to the known pilumnid zoeas in Korea and adjacent waters. *Journal of Crustacean Biology*, 23:341-351. [https://doi.org/10.1651/0278-0372\(2003\)023\[0341:ZDOHCD\]2.0.CO;2](https://doi.org/10.1651/0278-0372(2003)023[0341:ZDOHCD]2.0.CO;2)
- Lee DH, 1993. Larval development *Macromedaeus distinguendus* (De Haan, 1835) and *Pilumnopeus makiana* (Rathbun, 1929) (Brachyura, Xanthidae) reared in the laboratory. MS thesis, Pusan National University, Busan, Korea, pp. 1-53.
- Lee SH, 2020. The first zoeal stage of *Echinoecus nipponicus* (Decapoda: Pilumnidae: Eumedoninae) hatched in the laboratory. *Animal Systematics, Evolution and Diversity*, 36:304-308. <https://doi.org/10.5635/ASED.2020.36.4.030>
- Lee SH, Ko HS, 2009. Zoeal stages of *Harrovia japonica* (Decapoda: Brachyura: Pilumnidae) with a key to the known eumedoninid zoeae from the Indo-Pacific. *Animal Cells and Systems*, 13:213-222. <https://doi.org/10.1080/19768354.2009.9647213>
- Lee SK, Lee C, Lee SH, Lee SH, Song SJ, Khim JS, 2023. First report on occurrence and distributions of *Typhlocarcinops decrescens* Rathbun, 1914 and *Quadrella reticulata* Alcock, 1898, through long-term ecological surveys of the Korea marine and costal National Park in 2015-2020. *Regional Studies in Marine Science*, 59:102808. <https://doi.org/10.1016/j.rsma.2022.102808>
- Lee SK, Lee C, Noh J, Song SJ, Khim JS, 2021. First comprehensive ecological checklist of Brachyura in Korea: 1879-2020. *Marine Pollution Bulletin*, 171:112742. <https://doi.org/10.1016/j.marpolbul.2021.112742>
- Rice AL, 1980. Crab zoeal morphology and its bearing on the classification of the Brachyura. *Transactions of the Zoological Society of London*, 35:271-372. <https://doi.org/10.1111/j.1096-3642.1980.tb00060.x>
- Souza ASD, Costa RMD, Abrunhosa FA, 2013. Comparative morphology of the first zoea of twelve brachyuran species (Crustacea: Decapoda) from the Amazon region. *Zoologia Curitiba*, 30:273-290. <https://doi.org/10.1590/S1984-46702013000300004>
- WoRMS, 2024. *Zehntneriana* Ng and Takeda, 2010 [Internet]. *World Register of Marine Species*, Accessed 28 Nov 2024, <<https://www.marinespecies.org/aphia.php?p=taxdetails&id=557872>>.

Received December 3, 2024
 Revised December 16, 2024
 Accepted December 30, 2024